

**SAMPLE FILE FOR A THESIS WITH THE
'PITTED' CLASS**

by

I. M. Student

Previous degree, institution, year

Submitted to the Graduate Faculty of
the Department of Mathematics in partial fulfillment
of the requirements for the degree of

Doctor of Philosophy

University of Pittsburgh

2006

UNIVERSITY OF PITTSBURGH
MATHEMATICS DEPARTMENT

This dissertation was presented

by

I. M. Student

It was defended on

July 20th 1967

and approved by

Chair's name, Departmental Affiliation

Second member's name, Dept. Aff.

Third member's name, Dept. Aff.

Dissertation Advisors: Chair's name, Departmental Affiliation,

Second advisor, Dept. Aff.

SAMPLE FILE FOR A THESIS WITH THE ‘PITTETD’ CLASS

I. M. Student, PhD

University of Pittsburgh, 2006

This document is a sample file for the creation of ETD’s at Pitt through \LaTeX .

TABLE OF CONTENTS

1.0 SAMPLE MATHEMATICS AND TEXT	2
1.1 In-line and Displayed Mathematics	2
1.2 Mathematics in Section Heads $\int_{\alpha}^{\beta} \ln t dt$	3
1.3 Theorems, Lemmata, and Other Theorem-like Environments	3
BIBLIOGRAPHY	5
APPENDIX. TITLE OF APPENDIX	6
.1 First section of Appendix	6

1.0 SAMPLE MATHEMATICS AND TEXT

1.1 IN-LINE AND DISPLAYED MATHEMATICS

The expression $\sum_{i=1}^{\infty} a_i$ is in-line mathematics, while the numbered equation

$$\sum_{i=1}^{\infty} a_i \tag{1.1}$$

is displayed and automatically numbered as equation 1.1.

Let H be a Hilbert space, C be a closed bounded convex subset of H , T a non-expansive self map of C . Suppose that as $n \rightarrow \infty$, $a_{n,k} \rightarrow 0$ for each k , and $\gamma_n = \sum_{k=0}^{\infty} (a_{n,k+1} - a_{n,k})^+ \rightarrow 0$. Then for each x in C , $A_n x = \sum_{k=0}^{\infty} a_{n,k} T^k x$ converges weakly to a fixed point of T .

Two sets of L^AT_EX parameters govern mathematical displays.¹ The spacing above and below a display depends on whether the lines above or below are short or long, as shown in the following examples.

A short line above:

$$x^2 + y^2 = z^2$$

and a short line below.

A long line above may depend on your margins

$$\sin^2 \theta + \cos^2 \theta = 1$$

as will a long line below. This line is long enough to illustrate the spacing for mathematical displays, regardless of the margins.

¹L^AT_EX automatically selects the spacing depending on the surrounding line lengths.

1.2 MATHEMATICS IN SECTION HEADS $\int_{\alpha}^{\beta} \text{LN TDT}$

Mathematics can appear in section heads. Note that mathematics in section heads may cause difficulties in typesetting styles with running headers or table of contents entries.

1.3 THEOREMS, LEMMATA, AND OTHER THEOREM-LIKE ENVIRONMENTS

A number of theorem-like environments is available. The following lemma is a well-known fact on differentiation of asymptotic expansions of analytic functions.

Lemma 1. *Let $f(z)$ be an analytic function in \mathbb{C}_+ . If $f(z)$ admits the representation*

$$f(z) = a_0 + \frac{a_1}{z} + o\left(\frac{1}{z}\right),$$

for $z \rightarrow \infty$ inside a cone $\Gamma_{\varepsilon} = \{z \in \mathbb{C}_+ : 0 < \varepsilon \leq \arg z \leq \pi - \varepsilon\}$ then

$$a_1 = -\lim_{z \rightarrow \infty, z \in \Gamma_{\varepsilon}} z^2 f'(z), \quad (1.2)$$

Proof. Change z for $1/z$. Then $\Gamma_{\varepsilon} \rightarrow \bar{\Gamma}_{\varepsilon} = \{z \in \mathbb{C}_- : \bar{z} \in \Gamma_{\varepsilon}\}$ and

$$f(1/z) = a_0 + a_1 z + o(z). \quad (1.3)$$

Fix $z \in \bar{\Gamma}_{\varepsilon}$, and let $C_r(z) = \{\lambda \in \mathbb{C}_- : |\lambda - z| = r\}$ be a circle with radius $r = |z| \sin \varepsilon/2$. It follows from (1.3) that

$$\frac{1}{2\pi i} \int_{C_r(z)} \frac{f(\lambda) d\lambda}{(\lambda - z)^2} = \sum_{m=0}^1 a_m \frac{1}{2\pi i} \int_{C_r(z)} \frac{(\lambda - z_0)^m d\lambda}{(\lambda - z)^2} + R(z), \quad (1.4)$$

where for the remainder $R(z)$ we have

$$\begin{aligned} |R(z)| &\leq r^{-1} \max_{\lambda \in C_r(z)} o(|z|) = r^{-1} \max_{\lambda \in C_r(z)} |\lambda| \cdot O(|z| + r) \\ &= \frac{|z| + r}{r} \cdot O(|z| + r) = \frac{1 + \sin \varepsilon}{\sin \varepsilon} \cdot O(|z|). \end{aligned}$$

Therefore $R(z) \rightarrow 0$ as $z \rightarrow \infty, z \in \bar{\Gamma}_{\varepsilon/2}$, and hence by the Cauchy theorem (1.4) implies

$$\frac{d}{dz} f(1/z) = a_1 + R(z) \rightarrow a_1, \text{ as } z \rightarrow \infty, z \in \bar{\Gamma}_{\varepsilon/2},$$

that implies (1.2) by substituting $1/z$ back for z . □

Following is a short bibliography. It has no relationship to the previous text, but can be used to show sample citations such as [4] and [6]. This typesetting style places each citation inside square brackets. If you want multiple citations to appear in a single set of square brackets you must type all of the citation keys inside a single citation, separating each with a comma. Here is an example: [2, 3, 4].

Head	Head	Head
entry	entry	entry
entry	entry	entry
entry	entry	entry

Table 1: Sample table.

BIBLIOGRAPHY

- [1] American Petroleum Institute, Technical Data Book - Petroleum Refining, 5th edition, 1992
- [2] Harstad, K. and Bellan, J., "Isolated fluid oxygen drop behavior in fluid hydrogen at rocket chamber pressures", *Int. J. Heat Mass Transfer*, 1998a, **41**, 3537-3550
- [3] Harstad, K. and Bellan, J., "The Lewis number under supercritical conditions", *Int. J. Heat Mass Transfer*, in print
- [4] Hirshfelder, J. O., Curtis, C. F. and Bird, R. B., *Molecular Theory of Gases and Liquids*, John Wiley and Sons, Inc., 1964
- [5] Prausnitz, J., Lichtenthaler, R. and de Azevedo, E., *Molecular thermodynamics for fluid-phase equilibrium*, Prentice -Hall, Inc., 1986
- [6] Reid, R. C., Prausnitz, J. M. and Polling, B. E., *The Properties of Gases and Liquids*, 4th Edition, McGraw-Hill Book Company, 1987

APPENDIX

TITLE OF APPENDIX

.1 FIRST SECTION OF APPENDIX

This is the Appendix.